

# POSITIVE SOLUTIONS OF A LOGISTIC EQUATION ON UNBOUNDED INTERVALS

MA LI AND XU XINGWANG

ABSTRACT. In this paper, we study the existence of positive solutions of a one-parameter family of logistic equations on  $R_+$  or on  $R$ . These equations are stationary versions of the Fisher equations and the KPP equations. We also study the blow up region of a sequence of the solutions when the parameter approaches a critical value and the nonexistence of positive solutions beyond the critical value. We use the direct method and the sub and super solution method.

## 1. INTRODUCTION

As pointed out in the book written by G. Buttazzo, M. Giaguinta, and S. Hildebrandt [1], one dimensional variational problems deserve special attention. In fact, this kind of problems have their own characters. Sometimes, higher dimensional variational problems can be reduced to one dimensional ones.

In this work, we study the existence of positive solutions of the following one-parameter family of logistic equations  $(1)_a$ :

$$u'' + af(x)u - b(x)u^p = 0 \quad \text{in } I,$$

where  $I = (0, \infty)$  the half real line  $R_+$  or the whole real line  $R$ ,  $p > 1$ ,  $a \in R$  is the parameter,  $f(x)$  is a given  $C^1$  function on  $I$ , which is positive somewhere, and  $b(x)$  is a given non-negative smooth function on  $I$ . These equations are of interest in mathematical biology. Our equations appear also in Ginzburg-Landau theory of superconductivity [21]. Such equations were also known as the stationary version of the Fisher equation and KPP equation [20], and they were considered by Kazdan-Warner [6], Ouyang [7], and de Pino [8] in compact manifolds of higher dimensions and Du and Huang [10] in compact domains in the euclidean space  $R^n$ . Remarkably, Ouyang [7] proved a conjecture posed by Kazdan and Warner in [6]. Such problems have also been considered by Li-Tam-Yang [12], Afrouti-Brown [20] and Du-Ma [18-19] on  $R^n$ . For more background material, please see

---

1991 *Mathematics Subject Classification.* 53C, 35J.

*Key words and phrases.* direct method, blow-up, positive solution.

The work is in final form and no version of it will be submitted for publication elsewhere. The work of Ma is partially supported by 973 project of China, a grant from Ministry of Education, and a scientific grant of Tsinghua University at Beijing. This paper has been accepted for publication in Proc.AMS(2001).

[20]. However, to the authors's best knowledge, there are very few results on  $(1)_a$  on  $R_+$  or on  $R$ .

These problems have a close relation with the principal eigenvalue problem (2):

$$u'' + \lambda f(x)u = 0 \quad \text{in } R.$$

It is proved by K.J.Brown, C.Cosner, and J.Fleckinger [4] that if  $\int_R f(x)dx > 0$ , then there does not exist a positive principal eigenvalue of (2). This result is also true in dimension two. For more results about principal eigenvalue problems, we refer to the articles [5],[3] and [2]. So we may expect that the study of  $(1)_a$  has some special properties in dimesions one and two. We will study logistic equations in dimension two in the near future.

For simplicity, we write the left side of  $(1)_a$  by the operator  $T_a(u)$ , and we state the results in the case where  $I = R_+$ . Since we will use the sub and super method to study  $(1)_a$  in the case when  $I = R$ , the result will be stated later in this article. We now assume that  $I = R_+$ , and we say that  $f(x)$  is in *class P* if  $f(x)$  is positive some where and satisfies the condition that

$$\max\{f(x), 0\} \leq p(x), \quad \text{on } R_+,$$

where  $p(x)$  is some positive function on  $R_+$  with

$$\int_{R_+} xp(x)dx < \infty.$$

With this assumption we can obtain a compactness result, which is of the type of Rellich-Kondrakov imbedding for Sobolev spaces.

We define the zero set

$$M_0 = \{x \in R_+; b(x) = 0\},$$

which will play an important role in the existence problem and the blow-up problem (see [6]).

Assume that  $M_0$  is a bounded open set in  $R$ . Let  $\lambda_* := \lambda_1(M_0)$  and  $\lambda_1$  denote the principal eigenvalues of the operator  $-u''$  with respect to the weighted quotient

$$Q(u, M) := \int_M |u'|^2 / \int_M f(x)u^2$$

where we assume  $\int_M f(x)u^2 > 0$ ;  $M = M_0$  and  $M = R_+$  respectively. We write  $\lambda_* = +\infty$  when  $M_0$  is empty. Note that  $\lambda_* \geq \lambda_1$ .

We assume that  $f$  is positive somewhere outside  $M_0$ . So we have  $\lambda_* > \lambda_1$ . Then we will use the direct method to prove the existence of a positive solution of the equation  $(1)_a$  when  $a \in (\lambda_1, \lambda_*)$ . The idea goes as follows. Define the space  $E$  as the completion of  $C_0^1(R_+)$  with respect to the norm  $\int_R |u'|^2$ . Define the functional

$$J(u) = \frac{1}{2} \int_{R_+} (|u'|^2 - af(x)u^2) + \frac{1}{p+1} \int_{R_+} b(x)|u|^{p+1}$$

on  $E$ .  $J(u) = +\infty$  if  $\int_{R_+} b(x)|u|^{p+1} - af(x)u^2 = +\infty$ . Note that by our inequality (\*) to be proved in the next section, the functional  $J$  is well-defined on  $E$ . It is easy to see that  $J$  is weakly lower semi-continuous on  $E$ . Assume  $a \in (\lambda_1, \lambda_*)$ . We then prove that the functional  $J$  is coercive on  $E$  and we get a minimizer of the functional  $J$  on  $E$ . We will show that this minimizer is a non-trivial positive solution of  $(1)_a$ .

We also study the blow up region of a sequence of the solutions when the parameter approaches the critical value  $\lambda_*$  and the nonexistence of positive solutions beyond the critical value. We use a variational method and a barrier method.

To state our result, we introduce some notations and concepts. We will write  $\|u\|$  as the norm of  $u \in H$  and  $\|u\|_q$  as the norm of  $u \in L^q(R_+)$  where  $q > 1$ . We will write

$$L^2(R_+, p) = L^2(R_+, p dx).$$

We will sometimes write  $M := R_+$ .

Our main result is

**Theorem 1.** *Given a positive bounded function  $f(x)$  of class  $P$  and a non-trivial non-negative smooth function  $b$  on  $R_+$ . Assume that  $f$  is positive somewhere outside the zero set  $M_0$  of the function  $b$ . Then the following three assertions are true:*

- (i) *When  $a \in (\lambda_1, \lambda_*)$ , there is a unique positive smooth solution  $u_a \in H$  of  $(1)_a$ .*
- (ii) *Assume  $\lambda_* < +\infty$ . When  $a \rightarrow \lambda_*^-$ , we have that  $u_a \rightarrow +\infty$  on any compact set  $K \subset \subset M_0$ .*
- (iii) *Assume  $\lambda_* < +\infty$ . When  $a \geq \lambda_*$ , no positive  $E$ -solutions of  $(1)_a$  exist.*

We remark that we can extend the existence part of the result above to the following system:

$$W_j'' + aF_j(x)W_j - |W|^{p-1}W_j = 0, \quad \text{on } R_+,$$

with  $W_j(0) = 0$  for  $j = 1, \dots, N$ . Here we write  $W = (W_1, \dots, W_N)$  and assume that all the coefficients  $F_j(x)$  are dominated by functions in class  $(P)$ . Because the formulation of the result is straightforward, we omit it.

The rest of this paper is organized as follows. In section two, we set up some inequalities and we prove a compactness result. We also discuss the existence of principal eigenvalue and eigen-function problem (2) on  $R_+$ . Uniqueness and existence of positive  $E$ -solutions for  $(1)_a$  will be proved in section three. In section four and in section five, blow-up behavior of a solution sequence and non-existence of positive solutions will be presented. In the final section, we study the existence and uniqueness of positive solutions of  $(1)_a$  in the case when  $I = R$ .

## 2. COMPACTNESS LEMMATA

In this section we will set up some useful calculus results, which are easy to prove, but they are not easy to find in literatures (see [1] for more references). Hence we give full proofs. We point out that our compactness result is new.

**2.1. The case when  $I = R_+$ .** Assume that  $u \in C_0^1(R_+)$  with

$$\int_{R_+} |u'|^2 dx \leq C^2$$

where  $C$  is a positive constant. Then, for any  $x, y \in R_+$ , we have

$$|u(x) - u(y)| \leq \int_{[x,y]} |u'| dx.$$

Using Cauchy-Schwartz inequality we find

$$|u(x) - u(y)| \leq C|x - y|^{1/2}.$$

In particular, since  $u(0) = 0$ , we have

$$|u(x)| \leq C|x|^{1/2}.$$

Given a positive function  $p = p(x)$  on  $R_+$  satisfying

$$\int_{R_+} p(x)x dx < +\infty.$$

Then we have, for  $R \geq 0$ , that

$$\int_{(R,+\infty)} p(x)u^2 dx \leq C \int_{(R,+\infty)} p(x)x dx. \quad (*)$$

Define  $E$  as the completion of  $C_0^1(R_+)$  in the norm

$$|u| = \left( \int_{R_+} |u'|^2 dx \right)^{1/2}.$$

It is clear that the inequality (\*) is also true for every  $u \in E$ .

Let

$$L^2(R_+, p dx) = \{u \in L_{loc}^1(R_+); \int_{R_+} pu^2 < +\infty\}$$

Then we have the following compactness result.

**Proposition 2.** *The imbedding from  $E$  to  $L^2(R_+, p dx)$  is bounded and compact.*

*Proof.* The boundedness of the imbedding is clear from the inequality (\*).

Now let  $\{u_k\} \subset E$  with

$$\int_{R_+} |u'_k|^2 dx \leq C^2.$$

Then we have, for every  $k$  and any  $x, y \in R_+$ ,

$$|u_k(x) - u_k(y)| \leq C|x - y|^{1/2}.$$

In particular, take  $y = 0$  and we have

$$|u_k(x)| \leq C|x|^{1/2}.$$

Given  $\epsilon > 0$ . By inequality (\*) we can choose  $R > 1$  sufficient large such that for every  $k$  we have

$$\int_{[R,+\infty)} pu_k^2 \leq C \int_{[R,+\infty)} xp(x)dx < \epsilon/2.$$

By this and the Azela-Ascoli theorem we can find a subsequence, still denoted by  $\{u_k\}$ , such that it converges in  $C_{loc}^0(R_+)$ , and hence in  $L^2(R_+, p dx)$ . This completes the proof of compactness part of the theorem.  $\square$

Note that we can not handle the case when  $R_+$  is replaced by the entire real line  $R^1$  without further restriction. But we have another inequality, which will be introduced below. We remark that the inequality below can not be used to prove a similar compactness result as above.

Using the compactness result, one readily gets the following result on the existence of principal eigenvalue and function.

**Theorem 3.** *Assume the function  $f$  is positive somewhere on  $R_+$ , which is of the class (P). Then there exist a positive function  $V$  in  $E$  and a positive number  $\mu$  such that*

$$V'' + \mu f(x)V = 0, \quad \text{on } R_+$$

**2.2. The case when  $I = R$ .** Assume that  $p = p(x)$  is a positive function on the whole real line  $R^1$ . Now we define a Hilbert space  $E'$ , which is the completion of the space  $C_0^1(R^1)$  according to the Dirichlet integral. As before, we can define the space  $L^2(R, p dx)$ .

**Definition** *We call that a positive function  $p$  satisfies the condition (AB) if there are two positive constants  $A$  and  $B$  such that for any  $x \in R$ ,*

$$p(x) \leq \frac{A}{(|x| + B)^2}.$$

Although the following result will not be used later in this work, we believe that it is a useful tool in the analysis of solutions of ordinary differential equations.

**Proposition 4.** *Assume that the positive function  $p$  on  $R$  satisfies the condition (AB). Then there is a positive constant  $C$  such that for every  $u \in E'$ , it holds*

$$\int_R p(x)u^2 dx \leq \frac{4A}{B}u(0)^2 + C \int_R |u'|^2 dx.$$

We should mention that this inequality is a version of Hardy's inequality.

*Proof.* Without loss of generality, we may assume  $A = 1 = B$  and take  $u \in C_0^1(R)$ . Note that

$$\int_R \frac{1}{(|x|+1)^2} u^2 dx = \int_0^\infty \frac{u^2}{(x+1)^2} dx + \int_{-\infty}^0 \frac{u^2}{(x-1)^2} dx$$

Then by integration by part, we get

$$\int_0^\infty \frac{u^2}{(x+1)^2} dx = u(0)^2 + \int_0^\infty \frac{2uu'}{(x+1)} dx.$$

Similarly we have

$$\int_{-\infty}^0 \frac{u^2}{(x-1)^2} dx = u(0)^2 + \int_{-\infty}^0 \frac{2uu'}{(x-1)} dx.$$

Putting these two equalities together, we have

$$\int_R \frac{1}{(|x|+1)^2} u^2 dx = 2u(0)^2 + \int_R \frac{2\text{sgn}(x)uu'}{(|x|+1)} dx.$$

Note that by Cauchy-Schwartz inequality we find,

$$\int_R \frac{2|u||u'|}{(|x|+1)} dx \leq \frac{1}{2} \int_R \frac{1}{(|x|+1)^2} u^2 dx + 8 \int_R |u'|^2 dx.$$

Hence we get

$$\int_R \frac{1}{(|x|+1)^2} u^2 dx = 4u(0)^2 + 16 \int_R |u'|^2 dx.$$

□

### 3. AN APRIORI BOUND AND A COMPARISON LEMMA

In this section,  $a \in R$  is fixed. We write  $M = R_+$ .

As in the proof of Lemma 2.1 of [18] we have the following comparison lemma:

**Lemma 5.** *Assume  $\Omega$  is bounded open subset in  $M$ . Suppose  $u_1, u_2 \in C^2(\Omega)$  are positive in  $\Omega$ , and they satisfy  $T_a(u_1) \leq 0$  and  $T_a(u_2) \geq 0$  on  $\Omega$  and  $u_1 \geq u_2$  on  $\partial\Omega$ . Then  $u_1 \geq u_2$  on  $\Omega$ .*

We now use this result to obtain a local existence of of a boundary blow-up positive solution of  $(1)_a$  on a bounded interval  $G \subset M - M_0$ . This result may be of independent interest (see [18]).

**Proposition 6.** *Assume  $a \in R$ . There is at least one positive regular solution of the equation*

$$u'' + af(x)u - b(x)u^p = 0 \quad \text{in } G,$$

*with the boundary condition  $u = +\infty$  on  $\partial G$ .*

*Proof.* Fix an integer  $k \in \{1, 2, 3, \dots\}$ . As in the introduction we let

$$I(u) = \frac{1}{2} \int_G (|u'|^2 - af(x)u^2) + \frac{1}{p+1} \int_G b(x)|u|^{p+1}$$

defined on

$$E = \{u \in H^1(G); u|_{\partial G} = k\}.$$

It is easy to see that  $I$  is coercive on  $E$  (see section three for more difficult case) and there is a minimizer, which is denoted by  $u_k$ , of  $I$  on  $E$ .

By the comparison lemma above we have  $u_k \leq u_{k+1}$ , for  $k = 1, 2, \dots$ . By standard argument we can assume  $u_k \rightarrow U_a$  in  $C_{loc}^2(G)$ . This  $U_a$  is what we wanted.  $\square$

From the proof of the result above we can get the following

**Theorem 7.** *Assume that  $M_0$  is non-empty and the set  $M - M_0$  admits a bounded interval exhaustion, i.e., there is a bounded open subset sequence  $\{\Omega_j\}$  such that  $M - M_0 = \cup \Omega_j$ . Then for any  $a \in R$ , there is at least one positive smooth solution  $\bar{U}_a$  of  $T_a(u) = 0$  on  $M - M_0$  with the boundary blow-up condition  $\bar{U}_a = +\infty$  on  $M_0$ .*

We remark that when  $M_0$  is empty and  $a \leq 0$ , there may be no positive solutions of  $T_a(u) = 0$  on  $M$  at all (see [18]). One should also note that there is no contradiction between Theorem 1 (iii) with our Theorem 5 because we require  $T_a(u) = 0$  on the whole  $R_+$  in Theorem 1 (iii).

#### 4. EXISTENCE AND UNIQUENESS OF POSITIVE SOLUTIONS

We will discuss the uniqueness of positive  $E$ -solutions of  $(1)_a$  for  $a \in R$ . The proof of the following results are similar to the corresponding ones in [18], but for completeness, we give full proofs.

**Lemma 8.** *For  $a \in R$ , there is at most one positive  $E$ -solution to  $(1)_a$ .*

*Proof.* Assume  $u_1$  and  $u_2$  are two positive  $E$ -solutions of  $(1)_a$ . We assume  $u_1 \neq u_2$ , i.e., the set  $M_1 := \{x \in M; u_1(x) \neq u_2(x)\}$  has positive measure.

Let  $\epsilon_1 > \epsilon_2 > 0$  and denote

$$v_i = (u_i + \epsilon_i)^{-1}[(u_2 + \epsilon_2)^2 - (u_1 + \epsilon_1)^2]^+, \quad i = 1, 2$$

where  $[\cdot]^+$  is the positive part of  $[\cdot]$ . Then each  $v_i$  is an H-limit of a sequence in H with compact support. Hence, we have for  $i = 1, 2$ ,

$$-\int [u'_i v'_i - af u_i v_i] = \int b(x) u_i^p v_i.$$

By these we get

$$-\int [u'_2 v'_2 - u'_1 v'_1] = \int af(x)[u_1 v_1 - u_2 v_2] + \int b(x)(u_2^p v_2 - u_1^p v_1).$$

Note that the left side is

$$-\int (|u'_2 - \frac{u_2 + \epsilon_2}{u_1 + \epsilon_1} u'_1|^2 + |u'_1 - \frac{u_1 + \epsilon_1}{u_2 + \epsilon_2} u'_2|^2),$$

which is non-positive and goes to

$$-\int (|u'_2 - \frac{u_2}{u_1}u'_1|^2 + |u'_1 - \frac{u_1}{u_2}u'_2|^2) \leq 0.$$

The first term on the right goes to zero as  $\epsilon_1 \rightarrow 0$ . The second term on the right goes to

$$\int b(x)(u_2^{p-1} - u_1^{p-1})(u_2^2 - u_1^2),$$

which is non-negative by our assumption. Hence,  $M_1 \subset M_0$ . By all these we get that

$$\int_{M_0} (|u'_2 - \frac{u_2}{u_1}u'_1|^2 + |u'_1 - \frac{u_1}{u_2}u'_2|^2) = 0,$$

and

$$u_2 = cu_1, \quad \text{on } M_0$$

for some constant  $c \neq 1$ , a contradiction.  $\square$

In the rest of this section we assume that  $a \in (\lambda_1, \lambda_*)$ . We now prove part (i) of Theorem 1:

*Proof.* Recall that we assume  $M_0$  is bounded. It is clear that the functional  $J$  is weakly lower semi-continuous on  $E$  according to our compactness result. We claim that

$$J(u) \rightarrow +\infty \quad \text{as } \|u\| \rightarrow +\infty.$$

Assume not. Then we have sequence  $\{u_n\} \subset H$  such that  $J(u_n)$  is bounded above and  $\|u_n\| \rightarrow \infty$ . Then we have that  $d_n^2 := \int f(x)u_n^2 \rightarrow \infty$ ; for otherwise,  $\{\|u_n\|\}$  is bounded.

Set

$$\bar{u}_n = u_n/d_n.$$

Then we have  $\int f(x)\bar{u}_n^2 = 1$  and

$$\limsup \int_M |\bar{u}'_n|^2 - a + \frac{2}{(p+1)\|u_n\|_2^2} \int_M bu_n^{p+1} \leq 0.$$

By this, we may assume that  $\|\bar{u}_n\| + \int_M bu_n^{p+1}$  is uniformly bounded. Hence, we may assume further that  $\bar{u}_n \rightarrow \bar{u}$  weakly in  $E$ , almost everywhere, and strongly in  $L^2(R_+, p)$ . Then by Fatou's lemma, we get

$$\int_{R_+} b(x)\bar{u}^{p+1} = 0.$$

Hence,  $\bar{u} = 0$  on the set  $R_+ - M_0$ . Therefore, we can use the strong convergence on  $L^2(R_+, p)$  to get  $\int_{M_0} f(x)\bar{u}^2 = 1$  and

$$\int_{M_0} |\bar{u}'|^2 \leq a.$$

By this and the definition of  $\lambda_*$  we have

$$\lambda_* \leq a.$$

This is a contradiction with our hypothesis at the beginning.

Therefore, we have a minimizer  $u_0$ , which is a non-negative function, of the functional  $J$  on  $E$ . To prove that  $u_0$  is non-trivial, we take a large interval  $I_R$  with  $M_0 \subset I_R$  such that the first eigenvalue  $\lambda_1(I_R) - a \leq -\epsilon$  for some small  $\epsilon > 0$ . Let  $\phi_1$  be the eigenfunction of the eigenvalue  $\lambda_1(I_R)$  of the weighted quotient  $Q(u, I_R)$  and  $\int_{I_R} f(x)\phi^2 = 1$ . Then, for small  $t > 0$ ,

$$J(t\phi_1) = \frac{\lambda_1(I_R) - a}{2}t^2 + \frac{t^{p+1}}{p+1} \int_{I_R} b\phi^{p+1} < 0.$$

Note that  $J(0) = 0$ . Hence  $u_0$  is non-trivial and  $u_0 > 0$  everywhere on  $R_+$  by the strong maximum principle. The smoothness of  $u_0$  follows from the standard argument.

For the uniqueness of positive solutions, we use Lemma 8 above. Hence the proof is complete.  $\square$

**Remark:** By uniqueness of positive solution and the barrier method, we know immediately that, for  $x \in R_+$ ,  $a \rightarrow u_a(x)$  is increasing (for a proof, one may see [18]). This fact will be used later.

## 5. BLOW-UP OF THE SOLUTION SEQUENCE

In this section we assume  $\lambda_* < +\infty$ .

We study the behavior of the solution sequence when  $a_n \rightarrow \lambda_*^-$ . We assume  $a_n > \lambda_1$ . Set  $u_n = u_{a_n}$ .

We now prove Theorem 1(ii): First we note that

$$\lim_{n \rightarrow +\infty} \int_{R_+} f(x)u_n^2 = +\infty.$$

Otherwise, we can find a minimizer, which is the limit of  $u_n$ , of  $J$  as in Theorem 1(i), and this is impossible.

As in the proof of Theorem 1(i), we set

$$\bar{u}_n = u_n/d_n.$$

Then we find  $\bar{u} \in H$  with  $\bar{u} = 0$  on  $R_+ - M_0$ . Note that

$$\bar{u}_n'' + a_n f(x)\bar{u}_n = 0$$

on  $M_0$ . We can use the standard estimate [10] to conclude that the uniform convergence of  $\bar{u}_n$  to  $\bar{u}$  on  $K$ . This implies that  $u_n(x) \rightarrow +\infty$  on  $K$ .

## 6. NON-EXISTENCE OF POSITIVE SOLUTIONS

We prove in this section the third part of Theorem 1. That is, under the assumption  $\lambda_* < +\infty$ , there is no positive solution of  $(1)_a$  when  $a \geq \lambda_*$ .

Assume that  $a = \lambda_*$  and there is a positive solution, say  $u$ , of  $(1)_a$ . Let  $v$  be the  $\lambda_*$ -eigenfunction on  $M_0$  which satisfies the equation

$$u'' + af(x)u = 0, \quad \text{on } M_0$$

and the boundary condition  $v = 0$  on  $\partial M_0$ . Set  $\phi = v/u$ . Then by direct computation we find

$$\phi'' + 2\phi'(\log u)' = 0 \quad \text{on } M_0$$

and  $\phi = 0$  on  $\partial M_0$ . However, this gives an absurd result that  $\phi = 0$  on  $M_0$  by the standard maximum principle (see Theorem 3.1 in [10]). Therefore, there is no positive solution in this case.

In the case where  $a > \lambda_*$ . If there is a positive E-solution of  $(1)_a$ , say  $u_*$ . Then using the barrier method and the uniqueness result we find that  $u_a(x) \leq u_*(x)$  for all  $x \in M$  and  $a < \lambda_*$ . Therefore, there is no blow-up on  $M_0$  for any sequence of  $(u_a)$ , which is a contradiction with Theorem 1(ii). Hence, we get the conclusion of Theorem 1(iii).

We can give another proof for positive E-solutions when  $a > \lambda(M_0)$ . Take a smooth bounded domain  $G$  inside  $M_0$  such that  $\lambda_1(G) < a$ . Let  $w_1$  be the positive eigenfunction associated to  $\lambda_1(G)$ . Therefore, we can find a function  $u_0 \in E$  such that the number

$$\mu = \int_G (u_0' w_1' - a f(x) u_0 w_1)$$

is positive (see also [3]). Otherwise, we have

$$\int_{R_+} (u' w_1' - a f(x) u w_1) \leq 0,$$

for every  $u \in H$ . Hence, using  $-u$  as a test function, we have

$$\int_{R_+} (u' w_1' - a f(x) u w_1) = 0,$$

for every  $u \in H$ . Therefore,

$$-w_1'' - a f(x) w_1 = 0, \quad \text{on } R_+.$$

It is absurd with the definition of  $w_1$ .

Let

$$u_t = t w_1 + u_0$$

Then we compute  $J(u_t)$  and find

$$J(u_t) = (\lambda_1(G) - a)t^2/2 + \mu t + d$$

where

$$d = \frac{1}{p+1} \int_{R_+} b u_0^{p+1} + \frac{1}{2} \int_{R_+} (|u_0'|^2 - a f(x) u_0^2).$$

Hence,  $J(u_t) \rightarrow -\infty$  as  $t \rightarrow -\infty$ , which is a contradiction with the coercivity of  $J$ . So we conclude the result when  $a > \lambda_*$ .

## 7. THE REAL LINE CASE

In this section we discuss the equations  $(1)_a$  on the real line  $R$ . We point out that the method used here can not work for the case when  $b(x) = 0$  somewhere.

We assume that  $f$  is a bounded integrable function on  $R$ , and  $b(x)$  is positive and uniformly bounded away from zero, that is, there is an uniform constant  $c > 0$  such that (e):

$$b(x) \geq c > 0, \quad \text{for } x \in R.$$

For example,  $b(x) = 2 + \sin(x)$  satisfies the condition (e) above.

We assume  $a \in (\lambda_1, \lambda_*)$ .

Set  $I_k = (-k, k)$  for  $k = 1, 2, \dots$ . Then as before, we define the functional

$$J_k(u) = \frac{1}{2} \int_{I_k} (|u'|^2 - af(x)u^2) + \frac{1}{p+1} \int_{I_k} b(x)|u|^{p+1}$$

on  $E_k := H_0^1(I_k)$ . Clearly, the functional is bounded below (see [1]) on  $E_k$ . We can easily get a minimizer  $u_k \in E_k$  such that

$$J_k(u_k) = \inf_{u \in E_k} J_k(u),$$

and therefore we get a family of positive point-wise increasing functions  $(u_k)$ , which satisfy the equations  $(1)_a$  with the Dirichlet boundary condition  $u_k(k) = 0 = u_k(-k)$ .

Using the maximum principle we can easily show that  $u_k$  is uniformly bounded on  $R$ . In fact, at the maximum point  $x_0 \in I_k$ , we have that  $u''(x_0) \leq 0$ . Hence, by using our equation  $(1)_a$  we find

$$af(x_0)u(x_0) = b(x_0)u(x_0)^p - u''(x_0) \geq cu(x_0)^p$$

and  $u(x_0) \leq [aD/c]^{1/(p-1)}$ , where  $D = \sup_{x \in R} f_+(x)$ . According to our assumption on  $p$ , we know that  $u_k$  is uniformly bounded in  $L^2(R, pdx)$ .

Choose  $\xi \in C_0^1(I_1)$  and we have

$$J_k(u_k) \leq J_1(\xi)$$

for every  $k = 1, 2, \dots$ . Using this bound and the uniform boundedness of the sequence  $(u_k)$  we obtain an uniform bound of  $\int_R |u_k'|^2 dx$ . In fact, multiplying both sides of the equation (1) for  $u_k$  by  $u_k$  and integrating by part, we know that the integrals

$$\int_R u_k^{p+1} dx + \int_R |u_k'|^2 dx$$

are uniformly bounded. Therefore, we can pass to a subsequence, still denoted by  $(u_k)$ , which converges in  $C_{loc}^2(R)$  to a positive solution  $U$  of  $(1)_a$  on  $R$ . And it is easy to see that

$$\int_R |U'|^2 dx + \int_R U^{p+1} dx < +\infty$$

and  $U$  is in  $L^2(R, p dx)$ . Hence by standard estimate (see [18]), we have proven that this  $U$  is in  $E'$ . That is to say,  $U(x) \rightarrow 0$  uniformly as  $|x| \rightarrow \infty$ . As before, we can prove the uniqueness of the  $E'$ -solution.

We point out that for the case when  $a \in (0, \lambda_1]$ , we can easily show by the maximum principle that there is no positive  $E'$ -solution of  $(1)_a$ .

In summary, we have

**Theorem 9.** *Assume  $f$  is a bounded, positive somewhere, integrable function on  $R$ . Assume that the positive function  $b(x)$  satisfies (e) above. Then (i) for  $a \in (0, \lambda_1]$  there is no positive  $E'$ -solution of  $(1)_a$ ; (ii) for any  $a \in (\lambda_1, +\infty)$  there is a positive  $E'$ -solution of  $(1)_a$ , and this solution is unique in  $E'$ .*

We remark that some extension of Theorem 9 to high dimension has been obtained by Du and Ma [18].

#### REFERENCES

- [1] G. Buttazzo, M. Giaquinta, and S. Hildebrandt, *One-dimensional variational problems*, Oxford scientific publishing, Clarendon press, Oxford, 1998.
- [2] Z.R. Jin, *Principal eigenvalues with indefinite weight functions*, Trans. AMS, 349(1997)1945-1959.
- [3] J. Lopes-Gomez and J.C. Sabina de Lis, *First variation of principal eigenvalues*, J. Diff. Equa., 148(1998)47-64.
- [4] K.J. Brown, C. Cosner, and J. Fleckinger, *Principal eigenvalues for problems with indefinite weight function on  $R^N$* , Proc. AMS, 109(1990)147-155
- [5] W. Allegretto, *Principle eigenvalues for indefinite weight elliptic problems in  $R^N$* , Proc. AMS, 116(1992)701-706.
- [6] J. Kazdan-Warner, *Scalar curvature and conformal deformation of Riemannian structure*, J. Diff. Geom., 10(1985)113-134.
- [7] T. Ouyang, *On the positive solutions of semilinear equation  $\Delta u + \lambda u - hu^p = 0$  on the compact manifolds*, Trans. AMS, 331(1992)503-527.
- [8] de Pino, *Positive solutions of a semilinear elliptic equations on a compact manifold*, Nonlinear Analysis, 22(1994)1423-1430.
- [9] A. Edelson and A. Rumbos, *Linear and semilinear eigenvalue problems in  $R^n$* , Comm. Partial. Diff. Equations, 18, 215-240(1993).
- [10] Y. Du, Q. Huang, *Blow-up solutions for a class of semilinear elliptic and parabolic equations*, SIAM J. math. Anal., 31(2000)1-15..
- [11] L. Ma, *Conformal deformations on a noncompact riemannian manifold*, Math. Ann., 295(1993)75-80.
- [12] P. Li, L.F. Tam, D. Yang, *On the elliptic equation  $\Delta u + ku - Ku^p = 0$  on complete Riemannian manifolds and their geometric applications*, Trans. AMS, 350(1998)1045-1078.
- [13] W.M. Ni, *on the elliptic equation  $\Delta u + K(x)u^{(n+2)/(n-2)} = 0$ , its generalizations, and applications in geometry*, Indiana Univ. Math. J., 31(1982)495-529.
- [14] P. Aviles, R. McOwen, *conformal deformation to constant negative scalar curvature on non-compact Riemannian manifolds*, J. Diff. Geom., 27(1988)225-239.
- [15] Gilbarg and N.S. Trudinger, *Elliptic Partial Differential Equation of Second Order*, Springer-Verlag, 1977.
- [16] F.H. Lin, *on the elliptic equations  $D_i[A_{ij}D_j U] - k(x)U + K(x)U^p = 0$* , Proc. AMS, 95(1985)219-226.
- [17] M. Marcus and L. Veron, *Uniqueness and asymptotic behavior of solutions with boundary blow-up for a class of nonlinear elliptic equations*, Anal. Nonlinearie, 14(1997)237-274.
- [18] Y.H. Du and L. Ma, *Positive solutions of an elliptic partial differential equation on  $R^n$* , preprint, 2000.
- [19] Y.H. Du and L. Ma, *Logistic type equations on  $R^n$  by a squeezing method involving boundary blow-up solutions*, preprint, 2000.

- [20] G.A.Afrouzi and J.W.Brown, *On a diffusive logistic equation*, JMAA, 225(1998)326-339.
- [21] E.N.Dancer and S.P.Hasting, *On the global bifurcation diagram for the one dimensional Ginzburg-Landau model of superconductivity*, Europian J. Applied Math., 11(2000)271-291

XU: MATHEMATICS DEPARTMENT, THE NATIONAL UNIVERSITY OF SINGAPORE, 10 KENT RIDGE CRESCENT, SINGAPORE 119260  
*E-mail address:* `matxuxw@math.nus.edu.sg`

MA: DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA  
*E-mail address:* `lma@math.tsinghua.edu.cn`